

Activity Description

This is a worksheet containing a variety of activities to reinforce the content delivered in the 'Hydrogen: Properties, Thermodynamics and Electrochemistry' presentation. The content is aimed for student aged 15–18.

It consists of:

- Activities aimed at consolidating knowledge from the presentation
- An answer sheet for the activities

Hydrogen Properties – Worksheet



What are some key properties of hydrogen?

What are the key energy properties of gaseous hydrogen?

How does the energy density differ in liquid hydrogen? Why?

What is Hess's Law?

Draw a Hess Cycle to calculate the missing enthalpy...



What is a spontaneous reaction?

How can you make a non-spontaneous reaction occur? Can you relate this to an electrolyser?

Hydrogen Properties – Worksheet Answers

What are some key properties of hydrogen?

- Colourless
 - Tasteless
 - Odourless
 - Low density
 - Low solubility
 - Highly reactive
 - Highly flammable in air
 - Lightest element
-

What are the key energy properties of gaseous hydrogen?

- Low boiling point of -253°C (20K)
 - Low melting point of -259°C (14K)
 - Hydrogen as a fuel has:
 - A low energy density per unit volume
 - A high energy density per unit weight
-


How does the energy density differ in liquid hydrogen? Why?

- Liquid hydrogen has a much greater energy density per unit volume than gaseous hydrogen
 - This is because the particles in a liquid are much closer together than the particles in a gas (they are more densely packed together), hence the energy density of liquid hydrogen is much greater per unit volume
-

What is Hess's Law?

Hess's Law – the enthalpy change for a chemical reaction is independent of the route taken

Define these fuel cell components...



$$\Delta H = -(\Delta H_{\text{vaporisation}}) + \Delta H_{\text{reaction}} = -(+0.90) + (-284.93) = -285.83\text{kJ/mol}$$

- The reaction enthalpy is therefore -285.83kJ/mol , so it is exothermic

What is a spontaneous reaction?

- A spontaneous reaction is a reaction that occurs on its own
- No continuous outside energy source is required
- The more negative the Gibbs Free Energy for the reaction is, the more spontaneous
- Gibbs Free Energy is temperature dependent, so a change in temperature can force a reaction to occur spontaneously

How can you make a non-spontaneous reaction occur? Can you relate this to an electrolyser?

- Provide a continuous outside energy source
- Gibbs Free Energy is temperature dependent, so a change in temperature can force a reaction to occur spontaneously
- In an electrolyser, a continuous current is applied
- This allows a non-spontaneous reaction to occur, such as splitting water into hydrogen and oxygen